

Dissipation-Assisted Quantum Information Processing with Trapped Ions

A. Bermudez,¹ T. Schaetz,² and M. B. Plenio¹

¹*Institut für Theoretische Physik, Albert-Einstein Alle 11, Universität Ulm, 89069 Ulm, Germany*

²*Albert-Ludwigs-Universität Freiburg, Physikalisches Institut, Hermann-Herder-Strasse 3, 79104 Freiburg, Germany*

We introduce a scheme to perform dissipation-assisted quantum information processing in ion traps considering realistic decoherence rates, for example, due to motional heating. By means of continuous sympathetic cooling, we overcome the trap heating, and show that the damped vibrational excitations can still be exploited to mediate coherent interactions as well as collective dissipative effects. We describe how to control their relative strength experimentally, allowing for protocols of coherent or dissipative generation of entanglement. This scheme can be scaled to larger ion registers for coherent or dissipative many-body quantum simulations.

Among the different platforms for quantum computation (QC) [1] and quantum simulations (QS) [2], laser-cooled atomic ions in radio-frequency traps stand out as excellent small-scale prototypes [3, 4]. In the quest for large-scale QS, the development of micro-fabricated surface traps [5] provides a route towards large and arbitrary trapped-ion lattices capable of simulating a variety of exotic many-body models. In combination with the ability of transporting ions between different zones of the planar microchips, this technology would also allow for scalable QC [6]. Nonetheless, its success depends crucially on our ability to overcome the impact of various sources of decoherence. Of primary importance is the electric field noise from the trap electrodes, which leads to the so-called anomalous heating [7]. This heating becomes critical for the miniaturized traps required for scalability, where the ions lie closer to the electrodes and are subjected to stronger fluctuating fields, and even their gradients. Therefore, the anomalous heating hampers the scalability of phonon-mediated QC [8], and spin- and phonon-based QS [9] in surface traps.

A strategy to minimize this heating is to cool the complete setup to cryogenic temperatures [10]. Another possibility is to clean the surfaces of the electrodes by laser ablation [11] or ion bombardment [12], which has been shown to reduce the motional heating rates significantly. Although these approaches are very promising, a substantial residual noise still exists, and it would be highly desirable to develop alternative schemes to mitigate its impact. We hereby propose to overcome the anomalous heating by *continuously* applying sympathetic laser cooling during the whole QC or QS protocol.

The technique of sympathetic cooling requires active laser cooling on a particular subset of ions in a Coulomb crystal, while relying on the Coulomb interaction to passively cool the remaining ions. In addition to its use for precision spectroscopy, this technique may overcome the motional heating for QC purposes [13]. In fact, sympathetic ground-state cooling [14] has been implemented between the sequential application of two-qubit gates to recover high fidelities [15]. For scalable approaches based on miniaturization, the increasing impact of the motional heating reduces the fidelities achieved in this way. Therefore, it would be useful to perform sympathetic cooling continuously during the whole protocol. So far, three different schemes have been proposed: (i) In the absence of fluctuating electric-field gradients, one can mediate the interactions by certain vibrational modes robust to the heating, while sympathetically cooling the remaining modes during




Figure 1. **Dissipation-assisted quantum information:** (a) Two-isotope artificial Coulomb crystal with ions residing in the individual minima of a surface trap. The laser-cooled ions (red dots) assist the collective coherent/dissipative dynamics of the effective spins of the physical ions (blue dots). Both isotopes may be stored in the same individual minima without affecting the resultant lattice geometry. (b) Proof-of-principle experiment with three ions in a conventional rf-trap. The red arrows correspond to a standing wave providing the Doppler cooling of the central ion, whereas the blue arrows lead to a running wave tuned to the axial red sideband of the outer ions.

the operation [13]. (ii) By using far-detuned state-dependent dipole forces with negligible motional excitations [16], two-qubit gates do not rely on the motional coherence, and can thus withstand a heating/cooling that is considerably slower than the gate speed. (iii) For ground-state cooled crystals, active or sympathetic resolved-sideband cooling may provide a dissipative force during the gate operation. For gates faster than the heating, the success rate and fidelity of these operations can be substantially improved [17].

We emphasize that none of the above requirements are met in state-of-the-art surface traps. (i) Since the ions lie close to the surface electrodes, electric-field gradients avoid the isolation of robust vibrational modes whose excitations can mediate the interaction. (ii-iii) Heating rates in room-temperature setups (of the order of 1 phonon/ms [18]) coincide with the speed of far-detuned [16] and dissipation-assisted [17] gates, compromising the success of these protocols. In this work, we propose a dissipation-assisted protocol based on an always-on sympathetic cooling, which provides a different approach to overcome the anomalous heating. We identify regimes where the sympathetically-cooled vibrational modes can be either used as mediators of coherent interactions, or exploited to tailor the collective dissipation for quantum information processing. Since we only require to remain in the Doppler-cooling regime, this proposal can be applied to larger registers for QS.

The model.— We consider an arbitrary array of $N_t = N_\sigma + N_\tau$ trapped ions in equilibrium positions $\{\mathbf{r}_i^\sigma, \mathbf{r}_i^\tau\}$ (Fig. 1(a)). A

fraction of them, N_τ , is laser cooled via a dipole-allowed transition $|g\rangle \leftrightarrow |e\rangle$ with frequency ω_0^τ and decay rate Γ_τ , providing the sympathetic cooling of the remaining ions, N_σ . Two hyperfine ground-states $\{|\uparrow\rangle, |\downarrow\rangle\}$ form the spins/qubits for QS/QC purposes, such that their decay rate is negligible, and their energy spacing is ω_0^σ ($\hbar = 1$). To minimize the action of the cooling laser on the spins [14], one may use beams tightly focused on the τ -species, or exploit two ion species/isotopes with sufficiently-different transition frequencies.

By tracing out the electromagnetic bath, we obtain the master equation $\dot{\rho} = -i[H, \rho] + \mathcal{D}(\rho)$ with the Hamiltonian $H = H_\tau + H_\sigma + H_{\text{ph}} + V_{\sigma, \text{ph}} + V_{\tau, \text{ph}}$, and dissipator $\mathcal{D}(\rho)$. Here,

$$H_\tau = \frac{1}{2} \sum_l \omega_0^\tau \tau_l^z, \quad H_\sigma = \frac{1}{2} \sum_i \omega_0^\sigma \sigma_i^z, \quad H_{\text{ph}} = \sum_n \omega_n b_n^\dagger b_n, \quad (1)$$

represent the laser-cooled ions, the spins, and the vibrational excitations, respectively. We have introduced $\tau_l^\pm = |e_l\rangle\langle e_l| - |g_l\rangle\langle g_l|$, $\sigma_i^\pm = |\uparrow_i\rangle\langle \uparrow_i| - |\downarrow_i\rangle\langle \downarrow_i|$, and the creation/annihilation operators $b_n^\dagger, b_n$ for a particular phonon branch [19] with frequencies ω_n . The spin-phonon coupling required for QC/QS is provided by a pair of laser beams in a Raman configuration [20], which drive a red-sideband interaction

$$V_{\sigma, \text{ph}} = \sum_{in} \mathcal{F}_{in}^\sigma \sigma_i^+ b_n e^{-i\omega_\sigma t} + \text{H.c.}, \quad \mathcal{F}_{in}^\sigma = \frac{i\Omega_\sigma}{2} \eta_n^\sigma \mathcal{M}_{in} e^{i\mathbf{k}_\sigma \cdot \mathbf{r}_i^\sigma}, \quad (2)$$

where $\sigma_i^\pm = |\uparrow_i\rangle\langle \uparrow_i| - |\downarrow_i\rangle\langle \downarrow_i|$. Here, $\Omega_\sigma > 0$ is the two-photon Rabi frequency, and $\omega_\sigma(\mathbf{k}_\sigma)$ is the Raman frequency (wavevector). The Lamb-Dicke parameter $\eta_n^\sigma = \mathbf{k}_\sigma \cdot \mathbf{e}_p / \sqrt{2m_\sigma \omega_n}$ describes the laser coupling to the n -th normal mode with displacements $\mathcal{M}_{in}$ along $\mathbf{e}_p$, where m_σ is the ion mass. This spin-phonon coupling arises from the dipole laser-ion interaction after expanding $\eta_n^\sigma \ll 1$, such that $\omega_\sigma \approx \omega_0^\sigma - \omega_n$, and setting $\Omega_\sigma \ll \omega_n$ to neglect the contribution of the carrier terms. Additionally, we laser cool the τ -species in the node of a standing wave which is red-detuned from the dipole-allowed transition

$$V_{\tau, \text{ph}} = \sum_{ln} \mathcal{F}_{ln}^\tau \tau_l^+ Q_n e^{-i\omega_\tau t} + \text{H.c.}, \quad \mathcal{F}_{ln}^\tau = \frac{\Omega_\tau}{2} \eta_n^\tau \mathcal{M}_{ln}. \quad (3)$$

Here, $\tau_l^\pm = |e_l\rangle\langle e_l| - |g_l\rangle\langle g_l|$, $Q_n = b_n + b_n^\dagger$, and the remaining parameters are defined as those below Eq. (2), but applied to this particular transition. The differences between Eqs. (2) and (3) are due to the different regimes $\Gamma_\sigma \ll \omega_n \ll \Gamma_\tau$, which do not allow us to resolve the sidebands of the dipole-allowed transition. Note also that the carrier term is exactly cancelled only at the node of the standing wave. Finally, the dissipation is described by $\mathcal{D}(\rho) = \mathcal{D}_0(\rho) + \mathcal{D}_1(\rho)$, where

$$\mathcal{D}_0(\rho) = \sum_l \frac{1}{2} \Gamma_\tau (\tau_l^- \rho \tau_l^+ - \tau_l^+ \tau_l^- \rho) + \text{H.c.},$$

is the usual dissipator for a two-level atom at rest, where we have considered that the typical ion distances lead to negligible collective effects in the spontaneous emission. Finally,

$$\mathcal{D}_1(\rho) = \sum_{lm} \frac{1}{2} \Gamma_{\tau, nm} \tau_l^- (Q_n \rho Q_m - Q_n Q_m \rho) \tau_l^+ + \text{H.c.},$$

describes the recoil effects, where we have introduced $\Gamma_{\tau, nm} = 2\Gamma_\tau \alpha_q \eta_n^\tau \eta_m^\tau \mathcal{M}_{ln} \mathcal{M}_{lm}$, and $\alpha_q = (1 + 3q^2)/10(1 + q^2)$ depends on the linear/circular laser polarization $q = 0, \pm 1$ [21].

In addition to the Doppler-regime condition $\omega_n \ll \Gamma_\tau$, we further impose that $\mathcal{F}_{ln}^\tau \ll \omega_n, \Omega_\tau$. In this case, the laser-cooled ions reach the steady state very fast, and can be integrated out from the dynamics [22]. This can be accomplished by a Schrieffer-Wolff transformation for open quantum systems [23], which leads to the following master equation

$$\dot{\mu} = -i[H_\sigma + H_{\text{ph}} + V_{\sigma, \text{ph}}, \mu] + \tilde{\mathcal{D}}(\mu), \quad (4)$$

where $\mu = \text{Tr}_\tau\{\rho\}$, and we have introduced the dissipator

$$\tilde{\mathcal{D}}(\mu) = \sum_n \{ \Gamma_n^+ (b_n^\dagger \mu b_n - b_n b_n^\dagger \mu) + \Gamma_n^- (b_n \mu b_n^\dagger - b_n^\dagger b_n \mu) \} + \text{H.c.}$$

Here, the effective rates are expressed as $\Gamma_n^\pm = S(\mp \omega_n)$, where $S(\omega_n) = \int_0^\infty dt' e^{i\omega_n t'} \langle F_n(t') F_n(0) \rangle_{\text{ss}}$ is the steady-state fluctuation spectrum of the sideband terms $F_n = \frac{1}{2} \sum_l \Omega_\tau \eta_n^\tau \mathcal{M}_{ln} (\tau_l^+ + \tau_l^-)$, evaluated by means of the quantum regression theorem [22]. At this level, we can introduce the effects of the anomalous heating by substituting $\Gamma_n^+ \rightarrow \Gamma_n^+ = S(-\omega_n) + \Gamma_n^{\text{ah}}$, where Γ_n^{ah} is the anomalous heating rate for each mode. The cooling rates and mean phonon numbers are

$$W_n = \text{Re}\{\Gamma_n^- - \Gamma_n^+\}, \quad \bar{n}_n = \text{Re}(\Gamma_n^+)/W_n \quad (5)$$

Therefore, we can overcome the anomalous heating by shaping the fluctuation spectrum of the laser cooling force $W_n > 0$.

The master equation (4) describes an array of spins coupled to a set of damped vibrational modes. The idea now is to use the quanta of these modes, i.e. phonons, as mediators of a coherent spin-spin interaction. However, in addition to the coherent dynamics, the phonons also provide an indirect coupling to the electromagnetic reservoir that leads to some collective dissipation on the spins. Our goal is to find suitable regimes where these collective effects allow for QC/QS.

Tailoring the collective Liouvillian.— To guide our search for the correct regime, note that the two-qubit gates for QC presently implemented in different laboratories [8] use nearly-resonant spin-dependent forces. These gates rely on the motional coherence to suppress the residual spin-phonon entanglement. Since the motional coherence is absent in our case, the only possibility would be to work in the far off-resonant regime for QC[16] or QS[24], where $|\mathcal{F}_{in}^\sigma| \ll |\delta_n| \ll \omega_n$, such that $\delta_n = \omega_\sigma - (\omega_0^\sigma - \omega_n)$. We identify below the additional conditions to tailor the collective coherent/dissipative phonon-mediated processes in presence of sympathetic cooling.

For the particular values considered below, the laser-cooling rate reaches values of $W_n \approx 10^{-2} \omega_n$. In this case, the sympathetic cooling is very strong, and the vibrations reach the steady-state very fast. Hence, we can use a second Schrieffer-Wolff transformation to integrate out the phonons from Eq. (4), and obtain the following effective Liouvillian

$$\dot{\mu}_\sigma = \mathcal{L}_{\text{eff}}(\mu_\sigma) = -i[H_{\text{eff}}, \mu_\sigma] + \mathcal{D}_{\text{eff}}(\mu_\sigma), \quad (6)$$

where $\mu_\sigma = \text{Tr}_{\text{ph}}\{\mu\}$. Here, the coherent Hamiltonian is

$$H_{\text{eff}} = \sum_{i>j} \left(J_{ij}^{\text{eff}} \sigma_i^+ \sigma_j^- + \text{H.c.} \right) + \sum_{in} \frac{1}{2} B_{in}^{\text{eff}} \sigma_i^z,$$

which contains the phonon-mediated interactions of strength J_{ij}^{eff} , and an effective Zeeman splitting B_{in}^{eff} . The coupling to the phonons also introduces an indirect dissipation via

$$\mathcal{D}^{\text{eff}}(\mu_\sigma) = \sum_{i,j} \Gamma_{ij}^{\text{eff}} (2\sigma_i^+ \mu_\sigma \sigma_j^- - \sigma_j^- \sigma_i^+ \mu_\sigma - \mu_\sigma \sigma_j^- \sigma_i^+) \\ + \sum_{i,j} (\Gamma_{ij}^{\text{eff}} + \Gamma_{ij}^{\text{eff}}) (2\sigma_i^- \mu_\sigma \sigma_j^+ - \sigma_j^+ \sigma_i^- \mu_\sigma - \mu_\sigma \sigma_j^+ \sigma_i^-),$$

where $\Gamma_{ij}^{\text{eff}}, \Gamma_{ij}^{\text{eff}}$ are the effective dissipation rates. To find the correct regime, we must compare the time-scales derived from

$$J_{ij}^{\text{eff}} = -\sum_n \frac{\mathcal{F}_{in}^\sigma (\mathcal{F}_{jn}^\sigma)^*}{\delta_n^2 + W_n^2} \tilde{\delta}_n, \quad B_{in}^{\text{eff}} = -\frac{\mathcal{F}_{in}^\sigma (\mathcal{F}_{in}^\sigma)^*}{\delta_n^2 + W_n^2} \tilde{\delta}_n (2\bar{n}_n + 1), \\ \Gamma_{ij}^{\text{eff}} = +\sum_n \frac{\mathcal{F}_{in}^\sigma (\mathcal{F}_{jn}^\sigma)^*}{\delta_n^2 + W_n^2} W_n, \quad \Gamma_{ij}^{\text{eff}} = \sum_n \frac{\mathcal{F}_{in}^\sigma (\mathcal{F}_{jn}^\sigma)^*}{\delta_n^2 + W_n^2} W_n \bar{n}_n,$$

where we have introduced $\tilde{\delta}_n = \delta_n + \text{Im}\{\Gamma_n^+ - (\Gamma_n^-)^*\}$, and $\bar{n}_n$ stands for the steady-state mean phonon number after the sympathetic cooling. From the above expression, it is clear that by tuning the ratio $\mathcal{R}_n = W_n(\bar{n}_n + 1)/|\tilde{\delta}_n|$, we can control whether the phonon-mediated interactions prevail over the dissipation $\mathcal{R}_n \ll 1$, or vice versa $\mathcal{R}_n \gg 1$. Below, we estimate the orders of magnitude for realistic values of an experimental setup, and describe how to use both regimes for QC/QS.

Coherent and dissipative generation of entanglement.— We consider the simplest scenario to test our scheme: a three-ion chain in a linear Paul trap (Fig. 1(b)). To use realistic parameters, we take $^{25}\text{Mg}^{25+}$ - $^{24}\text{Mg}^{24+}$ - $^{25}\text{Mg}^{25+}$ as a prototype example, although we emphasize that the scheme also works for other species. We set the axial trap frequency to $\omega_z/2\pi = 4.1$ MHz, which is possible by optimizing the trap voltages. The dipole-allowed transition $|g\rangle = |3S_{1/2}\rangle \leftrightarrow |e\rangle = |3P_{1/2}\rangle$ of $^{24}\text{Mg}^{24+}$, which is characterized by $\lambda_\tau \approx 280.3$ nm and $\Gamma_\tau/2\pi \approx 41.4$ MHz, shall be used for continuous sympathetic cooling. By applying an external magnetic field, we can encode the spins in a couple of Zeeman sublevels $|F, M\rangle$ of the ground-state manifold of $^{25}\text{Mg}^{25+}$, namely $|\uparrow\rangle = |2, 2\rangle$ and $|\downarrow\rangle = |3, 3\rangle$. This leads to a resonance frequency of $\omega_0^\sigma/2\pi \approx 1.79$ GHz, and a negligible decay rate of $\Gamma_\sigma/2\pi \approx 10^{-14}$ Hz. Finally, a pair of weaker laser beams drive the axial red-sideband transition through a far-detuned excited state in the $3P_{3/2}$ manifold of $^{25}\text{Mg}^{25+}$, such that the Lamb-Dicke parameter is $\eta_1^\sigma \approx 0.16$.

The particular isotopic mass ratio $m_\tau/m_\sigma \approx 0.96$ implies that the axial vibrational modes are almost unchanged with respect to the homogeneous chain, $\omega_n/2\pi \approx \{4.1, 7.1, 10.1\}$ MHz. In order to attain a wide range of possible values for the ratio $\mathcal{R}_n$, we tune the Raman lasers closer to the so-called egyptian mode, such that the detunings are $\delta_n/2\pi \in \{6.2, 3.2, 0.3\}$ MHz. Note that due to the large detuning from the remaining modes, the collective effects will be mediated by the egyptian mode. To sympathetically cool it, we place the cooling isotope at the middle of the chain, such that the standing-wave frequency is red-detuned $\omega_\tau = \omega_0^\tau + \Delta_\tau$ with $\Delta_\tau = -\Gamma_\tau/2$. According to Eq. (5), this leads to a steady-state mean phonon number $\bar{n}_3 = 0.65$ independent of the standing-wave Rabi frequency. Therefore, we can modify the Rabi frequency $0.2 \leq \Omega_\tau/\Gamma_\tau \leq 2$ in order to control the cooling rate




Figure 2. **Coherent/dissipative generation of entanglement:** (a) Coherent exchange dynamics (main panel) for the Liouvillian (4) specific to the egyptian mode (solid lines) and the effective description (6) (circles and squares). Bell-state fidelity $\mathcal{F} = |\langle \psi_B | \mu_\sigma | \psi_B \rangle|$ during a single SWAP (left lower panel), and optimization of the fidelity for different gate times t_B (right lower panel). Solid lines represent the complete Liouvillian (4), whereas the stars follow from the effective description (6). (b) Dissipative dynamics (upper panel) under Eq. (4) (solid lines) and Eq. (6) (circles and squares). Bell-state fidelity $\mathcal{F} = |\langle \phi_- | \mu_\sigma | \phi_- \rangle|$ at the steady state (lower panel).

W_3 , and thus the ratio $\mathcal{R}_3$, which shall allow us to explore regimes of either dominant collective dissipation or coherent interactions. We remark that the laser used for cooling $^{24}\text{Mg}^{24+}$ will be highly detuned from the cooling transition of $^{25}\text{Mg}^{25+}$ (i.e. $\Delta/2\pi \approx 2.7$ THz), such that the induced decay rate for the considered regime lies in $\Gamma_\tau(\Omega_\tau/\Delta)^2/(2\pi) \leq 10^{-2}$ Hz, and will not affect the dynamics. This laser only contributes with off-resonant ac-Stark shifts that shall be considered later on. We now explore two possible applications for this setup.

(a) Coherent generation of entanglement: Our goal is to use the coherent phonon-mediated interaction to generate entanglement between the $^{25}\text{Mg}^{25+}$ ions. By setting $\Omega_\sigma = 0.15\Gamma_\tau$, we obtain a cooling rate of $W_3/\omega_3 \approx 4.3 \cdot 10^{-3}$, such that $\mathcal{R}_3 \approx 7 \cdot 10^{-3}$, and the Hamiltonian part of the Liouvillian (6) thus dominates. Initializing the spin state in $|\psi_\sigma(0)\rangle = |\uparrow_1\downarrow_3\rangle$, and setting $\Omega_\sigma\eta_3^\sigma \approx 10W_3$, such that the distance between the ions is an integer multiple of the effective Raman wavelength, we obtain the Bell state $|\psi_B\rangle = \frac{1}{\sqrt{2}}(|\uparrow_1\downarrow_3\rangle - i|\downarrow_1\uparrow_3\rangle)$ for $t_f \approx 4$ ms (Fig. 2(a)). We have considered a heating rate $\Gamma_{\text{ah}} \approx 0.1$ phonon/ms for the rf trap. Even if the process is slower than the usual gates [8], it prevails over the phonon-mediated decoherence, and leads to fidelities above 98%.

(b) Dissipative generation of entanglement: A different possibility is to exploit the collective dissipation to generate entanglement in the steady state. The idea is to set $\Omega_\tau = 2\Gamma_\tau$, such that $W_3/\omega_3 \approx 0.03$ and $\mathcal{R}_3 \approx 2.3$, and the dissipative part of the Liouvillian (6) becomes dominating. In this regime, we can exploit a superradiant/subradiant phenomenon [25]. Considering the different decay channels $\Gamma_\pm^{\text{eff}} = \Gamma_{11}^{\text{eff}} \pm \Gamma_{12}^{\text{eff}}$ of the symmetric and antisymmetric Bell states $|\phi_\pm\rangle = \frac{1}{\sqrt{2}}(|\uparrow_1\downarrow_3\rangle \pm |\downarrow_1\uparrow_3\rangle)$, it is possible to set the ratio between the ions distance and the Raman wavelength, such that either $\Gamma_+^{\text{eff}} = 0$ or $\Gamma_-^{\text{eff}} = 0$. We note that this phononic superradiance goes beyond the possibilities of the pioneer experiments of pho-

tonic superradiance with trapped ions [26]. In particular, starting from $|\psi_\sigma(0)\rangle = |\uparrow_1\downarrow_3\rangle$ in the regime of $\Gamma_-^{\text{eff}} = 0$, we can generate entangled states dissipatively. After setting $\Omega_\sigma \eta_3^\sigma \approx W_3$, we obtain a decoherence-free steady-state Bell state $|\phi_-\rangle$ for $t \gg t_{ss} \approx 50 \mu\text{s}$ (Fig. 2(b)) with fidelities around 30%. We note that the ultimate limit of 50% cannot be achieved due to the thermal contribution to Eq. (6). However, schemes originally formulated for cavities [27], can also be applied in our trapped-ion setting to reach unit fidelities.

Note that the typical time-scales of coherent and dissipative processes are 0.1-10ms. In this regime, some other sources of noise may also become relevant. We show below that our protocol can be extended to make it robust against such noises.

Other sources of noise.— Fluctuating magnetic fields, laser intensities and thermal noise, lead to the dephasing term

$$H_n = \sum_i \frac{1}{2} (B_{in}^{\text{eff}} + F_i(t)) \sigma_i^z. \quad (7)$$

Here, B_{in}^{eff} introduces noise via thermal fluctuations over the steady-state mean phonon number, and $F_i(t)$ is a Markovian Gaussian random process that models the noise of external magnetic fields and uncompensated ac-Stark shifts, such as those provided by the cooling laser. This term leads to a dephasing rate of $\Gamma_d = \frac{1}{2} \int_0^\infty dt' \langle F_j(t') F_j \rangle_{\text{st}} = 1/2T_2$, where we have assumed a local noise with a short correlation time τ_c , and $\langle F_j(t') F_i \rangle_{\text{st}} = \frac{2\Gamma_d}{\tau_c} e^{-t'/\tau_c} \delta_{ji}$. Let us note that that typical decoherence times are of the order $T_2 \approx 1\text{-}10\text{ms}$ ($\Gamma_d \approx 0.05\text{-}0.5\text{kHz}$), and that for the regimes considered above, one obtains $B_{in}^{\text{eff}}/\bar{n}_n \approx 0.3\text{-}3\text{kHz}$. Comparing these values to the above time-scales ($\sim 0.1\text{-}10\text{ms}$), the need for a scheme to actively decouple from these noise sources becomes rather clear. A partial solution are the so-called clock states, such as $|\uparrow\rangle = |2, 1\rangle$ and $|\downarrow\rangle = |3, 1\rangle$ at a field of $B_0 = 213\text{G}$ for $^{25}\text{Mg}^+$, which become insensitive to linear Zeeman shifts. However, we still need to mitigate the other sources of dephasing by a different mechanism. As a byproduct, this decoupling mechanism yields also a tool to tailor the collective Liouvillian.

The standard approach for prolonging the coherence of a system consists of a sequence of pulses that refocus the induced dephasing. This technique is known as pulsed dynamical decoupling [28], and has already been applied to ions [29]. Another approach known as continuous dynamical decoupling, produces a similar effect by using continuous drivings, and has also been successfully applied to ions [30]. We apply a strong continuous driving resonant with the spins [31], such that the bare spin Hamiltonian (1) now reads

$$H_\sigma = \frac{1}{2} \sum_i \omega_\sigma \sigma_i^z + (\Omega_d \sigma_i^+ \cos(\omega_d t) + \text{H.c.}).$$

Here, the driving frequency is set to $\omega_d = \omega_\sigma$, and its Rabi frequency $|\Omega_d| \ll \omega_\sigma$. Note that the considered resonance frequencies lie in the microwave regime $\omega_\sigma/2\pi \approx 1\text{GHz}$, such that the driving can still be strong enough to fulfill

$$\{W_n, \tilde{\delta}_n\} \ll |\Omega_d|, \quad \Gamma_d, \tau_c^{-1} \ll |\Omega_d|. \quad (8)$$

In this regime, we apply a modified Schrieffer-Wolff transformation to obtain a new master equation $\dot{\mu}_\sigma = \tilde{\mathcal{L}}_{\text{eff}}(\mu_\sigma)$, where the collective Liouvillian is modified as follows

$$\tilde{\mathcal{L}}_{\text{eff}}(\mu_\sigma) = -i[\tilde{H}_{\text{eff}} + \tilde{H}_n, \mu_\sigma] + \tilde{\mathcal{D}}_{\text{eff}}(\mu_\sigma) + \tilde{\mathcal{N}}_n(\mu_\sigma). \quad (9)$$

The coherent Hamiltonian above describes a collection of spins interacting according to the so-called Ising model

$$\tilde{H}_{\text{eff}} = \sum_{i>j} \tilde{J}_{ij}^{\text{eff}} \sigma_i^x \sigma_j^x + \sum_{in} \frac{1}{2} \Omega_d \sigma_i^x,$$

whereas the phonon-mediated dissipation corresponds to a collective dephasing in the eigenbasis of the strong driving

$$\tilde{\mathcal{D}}_{\text{eff}}(\mu_\sigma) = \sum_{i,j} \tilde{\Gamma}_{ij}^{\text{eff}} (2\sigma_i^x \mu_\sigma \sigma_j^x - \sigma_j^x \sigma_i^x \mu_\sigma - \mu_\sigma \sigma_j^x \sigma_i^x).$$

Let us emphasize that the collective interaction strengths and dissipation rates, which are expressed as follows

$$\tilde{J}_{ij}^{\text{eff}} = -\sum_n \frac{\mathcal{F}_{in}^\sigma (\mathcal{F}_{jn}^\sigma)^*}{\tilde{\delta}_n^2 + W_n^2} \frac{\tilde{\delta}_n}{2}, \quad \tilde{\Gamma}_{ij}^{\text{eff}} = \sum_n \frac{\mathcal{F}_{in}^\sigma (\mathcal{F}_{jn}^\sigma)^*}{\tilde{\delta}_n^2 + W_n^2} \frac{W_n}{2} (\bar{n}_n + \frac{1}{2}),$$

lead to a very similar control parameter $\tilde{\mathcal{N}}_n = W_n(\bar{n}_n + \frac{1}{2})/|\tilde{\delta}_n|$. Accordingly, under the same assumptions considered above, we can interpolate between regimes where the coherent Ising interactions dominate over the collective Ising dissipation, or viceversa. In addition to the new range of possibilities offered by this collective Liouvillian, note that the noise terms in (7) contribute to the new Liouvillian (9) as follows

$$\tilde{H}_n = \sum_i \frac{1}{2} \tilde{\Omega}_d \sigma_i^x, \quad \tilde{\mathcal{N}}_n(\mu_\sigma) = \sum_i \tilde{\Gamma}_d (\sigma_i^x \mu_\sigma \sigma_i^x - \mu_\sigma),$$

where $\tilde{\Omega}_d = (B_{jn}^{\text{eff}} \tilde{\delta}_n \tau_c + 2\Gamma_d)/2\Omega_d \tau_c$, and $\tilde{\Gamma}_d = \Gamma_d/(\Omega_d \tau_c)^2$. According to the constraints (8), these noisy terms are clearly suppressed by a sufficiently strong driving. To be more specific, for noise correlation times on the order of $\tau_c = 10^{-2}T_2$ and detunings $\tilde{\delta}_n/2\pi \sim 0.1\text{-}1\text{MHz}$, it suffices to apply driving Rabi frequencies of $\Omega_d/2\pi \approx 10\text{MHz}$ to reduce both sources of noise by more than two orders of magnitude. Therefore, we are decoupling from the noise very efficiently, while preserving the collective part of the Liouvillian for QC and QS. In contrast to Fig. 2(a), we remark that the new Liouvillian (9) allows for the coherent generation of all four Bell states with similar fidelities and gate times. Besides, as a consequence of our control over the collective dephasing, any of these Bell states is made decoherence-free by tuning the ratio of the ion distance and the laser wavelength appropriately.

Many-body physics.— The collective Liouvillians in Eqs. (6) and (9) define our toolbox for dissipation-assisted QS. So far, we have considered quantum information processing by means of an isolated vibrational mode that mediates the collective dynamics. In the many-ion scenario, this would lead to long-range interactions that define a fully-connected spin network. For surface traps (Fig. 1(a)), the presence of magnetic frustration already leads to rich phase diagrams [32]. However, to achieve strongly-correlated models, it is better to work with a full vibrational branch of a small frequency width [24]. The small ratios $\mathcal{N}_n \sim 10^{-3}$ obtained for trapping frequencies $\omega/2\pi \approx 10\text{MHz}$, indicate that the coherent dynamics can be dominating also in this case, while minimizing the heating. This would allow for the QS of exotic models [33] with 3-body interactions or topological order in surface ion traps.

These conventional many-body QS focus on collective properties of the Hamiltonian while minimizing the influence of the environment. However, the dissipative dynamics may also lead to interesting many-body phenomena [34]. The possibility to control the relative strength of the coherent/dissipative parts in (6) and (9) is very appealing in this respect. We note that for intermediate driving strengths Ω_d , competing dissipative terms in (9) can lead to purely-dissipative quantum phase transitions.

Conclusions and Outlook.— We have proposed a scheme based on sympathetic cooling to overcome the anomalous

heating in surface traps, while allowing for QC and QS. In addition, the sympathetic cooling becomes a tool to tailor the collective effects of the Liouvillian. By controlling a single parameter, namely the laser-cooling power, we have shown how the Liouvillian interpolates between regimes of dominating coherent interactions/collective dissipation, both of which allow for generation of entanglement. Moreover, we have pointed out how this control can be exploited for coherent/dissipative many-body QS.

Acknowledgements.— This work was supported by PICC, DFG (SCHA973/1-6), and by the Alexander von Humboldt Foundation. A.B. thanks FIS2009-10061, and QITEMAD. We thank A. Albrecht and J. Almeida for useful discussions.

-
- [1] T. D. Ladd, F. Jelezko, R. Laflamme, Y. Nakamura, C. Monroe, and J. L. O'Brien, *Nature* **464**, 45 (2010).
 - [2] J. I. Cirac and P. Zoller, *Nat. Phys.* **8**, 264 (2012).
 - [3] R. Blatt and D. Wineland, *Nature* **453**, 1008 (2008).
 - [4] R. Blatt and C. F. Roos, *Nat. Phys.* **8**, 277 (2012).
 - [5] See D. J. Wineland and D. Leibfried, *Laser Phys. Lett.* **8**, 175 (2011) and references therein.
 - [6] D. Kielpinski, C. Monroe, and D. J. Wineland, *Nature* **417**, 709 (2002).
 - [7] Q. A. Turchette, D. Kielpinski, B. E. King, D. Leibfried, D. M. Meekhof, C. J. Myatt, M. A. Rowe, C. A. Sackett, C. S. Wood, W. M. Itano, C. Monroe, and D. J. Wineland, *Phys. Rev. A* **61**, 063418 (2000).
 - [8] See K.-A. Brickman Soderberg and C. Monroe, *Rep. Prog. Phys.* **73**, 036401(2010), and references therein.
 - [9] See Ch. Schneider, D. Porras, and T. Schaetz, *Rep. Prog. Phys.* **75**, 024401 (2012), and references therein.
 - [10] L. Deslauriers, S. Olmschenk, D. Stick, W. K. Hensinger, J. Sterk, and C. Monroe, *Phys. Rev. Lett.* **97**, 103007 (2006); J. Labaziewicz, Y. Ge, D. Leibbrandt, K. R. Brown, and I. L. Chuang, *Phys. Rev. Lett.* **100**, 013001 (2008).
 - [11] D. T. C. Allcock, L. Guidoni, T. P. Harty, C. J. Ballance, M. G. Blain, A. M. Steane, and D. M. Lucas, *New J. Phys.* **13**, 123023 (2011).
 - [12] D. A. Hite, Y. Colombe, A. C. Wilson, K. R. Brown, U. Warring, R. Jördens, J. D. Jost, D. P. Pappas, D. Leibfried, and D. J. Wineland, *Phys. Rev. Lett.* **109**, 103001 (2012).
 - [13] D. Kielpinski, B. E. King, C. J. Myatt, C. A. Sackett, Q. A. Turchette, W. M. Itano, C. Monroe, D. J. Wineland, and W. H. Zurek, *Phys. Rev. A* **61**, 032310 (2000); G. Morigi and H. Walther, *Eur. Phys. J. D* **13**, 261 (2001).
 - [14] H. Rohde, S. T. Gulde, C. F. Roos, P. A. Barton, D. Leibfried, J. Eschner, F. Schmidt-Kaler, R. Blatt, *J. Opt. B* **3**, S3 (2001); M. D. Barrett, B. DeMarco, T. Schaetz, V. Meyer, D. Leibfried, J. Britton, J. Chiaverini, W. M. Itano, B. Jelenkovic, J. D. Jost, C. Langer, T. Rosenband, and D. J. Wineland, *Phys. Rev. A* **68**, 042302 (2003); J. P. Home, M. J. McDonnell, D. J. Szwer, B. C. Keitch, D. M. Lucas, D. N. Stacey, and A. M. Steane, *Phys. Rev. A* **79**, 050305(R) (2009).
 - [15] J. P. Home, D. Hanneke, J. D. Jost, J. M. Amini, D. Leibfried, D. J. Wineland, *Science* **325**, 1227 (2009).
 - [16] A. Sørensen and K. Mølmer, *Phys. Rev. Lett.* **82**, 1971 (1999).
 - [17] A. Beige, *Phys. Rev. A* **67**, 020301(R) (2003); *ibid.* **69**, 012303 (2004).
 - [18] K.R. Brown, C. Ospelkaus, Y. Colombe, A. C. Wilson, D. Leibfried, and D. J. Wineland *Nature* **471**, 196 (2011); M. Harlander, R. Lechner, M. Brownnutt, R. Blatt, and W. Hänsel *Nature* **471**, 200 (2011).
 - [19] The scheme works for any vibrational branch of the crystal (e.g. axial and radial in a chain) provided that the Coulomb couplings and sympathetic cooling rates are of comparable magnitude.
 - [20] See D. J. Wineland, C. Monroe, W. M. Itano, D. Leibfried, B. E. King, and D. M. Meekhof, *J. Res. Natl. I. St. Tech.* **103**, 259 (1998), and references therein.
 - [21] J. Dalibard and C. Cohen-Tannoudji, *J. Phys. B: At. Mol. Phys.* **18**, 166 (1985); Y. Castin, H. Wallis, and J. Dalibard, *J. Opt. Soc. Am. B*, **6**, 2046 (1989).
 - [22] J. I. Cirac, R. Blatt, P. Zoller, and W. D. Phillips, *Phys. Rev. A* **46**, 2668 (1992).
 - [23] E. M. Kessler, *Phys. Rev. A* **86**, 012126 (2012).
 - [24] D. Porras and J. I. Cirac, *Phys. Rev. Lett.* **92**, 207901 (2004).
 - [25] R. H. Dicke, *Phys. Rev.* **93**, 99 (1954); R. H. Lemberg, *Phys. Rev. A* **2**, 889 (1970); Z. Ficek and R. Tanas, *Phys. Rep.* **372**, 369 (2002).
 - [26] R. G. DeVoe and R. G. Brewer, *Phys. Rev. Lett.* **76**, 2049 (1996).
 - [27] M. J. Kastoryano, F. Reiter, and A. S. Sørensen, *Phys. Rev. Lett.* **106**, 090502 (2011).
 - [28] E. L. Hahn, *Phys. Rev.* **80**, 580 (1950); L. Viola and S. Lloyd, *Phys. Rev. A* **58**, 2733 (1998).
 - [29] M. J. Biercuk, H. Uys, A. P. VanDevender, N. Shiga, W. M. Itano, and J. J. Bollinger, *Nature* **458**, 996 (2009); D. J. Szwer, S. C. Webster, A. M. Steane, and D. M. Lucas, *J. Phys. B* **44**, 025550 (2011).
 - [30] N. Timoney, I. Baumgart, M. Johanning, A. F. Varon, Ch. Wunderlich, M. B. Plenio, and A. Retzker, *Nature* **476**, 185 (2011); A. Noguchi, S. Haze, K. Toyoda, and S. Urabe, *Phys. Rev. Lett.* **108**, 060503 (2012).
 - [31] A. Bermudez, P. O. Schmidt, M. B. Plenio, and A. Retzker, *Phys. Rev. A* **85**, 040302(R) (2012).
 - [32] G.-D. Lin, C. Monroe, L.-M. Duan, *Phys. Rev. Lett.* **106**, 230402 (2011).
 - [33] A. Bermudez, D. Porras, and M. A. Martin-Delgado, *Phys. Rev. A* **79**, 060303(R) (2009); R. Schmied, J. H. Wesenberg, and D. Leibfried, *New J. Phys.* **13**, 115011 (2011).
 - [34] M.B. Plenio and S.F. Huelga, *Phys. Rev. Lett.* **88**, 197901 (2002); S.F. Huelga and M.B. Plenio, *Phys. Rev. Lett.* **98**, 170601 (2007); M. Müller, S. Diehl, G. Pupillo, P. Zoller, *Adv. At. Mol. Opt. Phys.* **61**, 1 (2012).